



Détection précoce de la dysfonction atriale gauche chez les patients hypertendus: Intérêt du Speckle Tracking

Early detection of left atrial dysfunction in hypertensive patients: Role of Speckle Tracking imaging

Karima Taamallah, Wael Yaakoubi, Abdeddayem Haggui, Nadhem Hajlaoui, Wafa Fehri

Hôpital Militaire Principal d'Instruction de Tunis / Faculté de médecine de Tunis

RÉSUMÉ

Introduction: L'hypertension est un problème de santé majeur dans le monde, elle est associée à une altération de la fonction de l'oreillette gauche (OG). L'analyse de la déformation myocardique à l'aide de l'échocardiographie s'avère un outil prometteur pour évaluer de la fonction auriculaire gauche.

Objectif : Evaluer les changements précoces du strain atrial gauche chez des patients souffrant d'hypertension.

Methodes : Le strain de l'OG a été étudié chez 109 patients hypertendus sans dilatation de l'OG et 50 témoins appariés selon l'âge. Des évaluations échocardiographiques bidimensionnelles et en speckle tracking ont été réalisées et les paramètres suivants ont été mesurés : le pic longitudinal du strain atrial gauche et la vitesse du strain pendant les périodes de la phase réservoir, de la phase conduite et de la contraction atriale gauche dans les vues à quatre et deux chambres. Le temps nécessaire pour atteindre le pic longitudinal du strain atrial gauche et la vitesse du strain ont été mesurés durant les trois phases de la fonction OG

Resultats : Le diamètre antéro-postérieur était normal, aucune différence entre les patients hypertendus et les témoins n'a été notée (34.35 mm \pm 4.91 vs 31.82 mm \pm 4.87, $p=0.16$). Les volumes maximum (41.78 \pm 10.29 vs 47 \pm 13.21, $p=0.01$), minimum (23.95 \pm 12.18 vs 16.94 \pm 7.91, $p=0.001$) de l'OG étaient plus élevés chez les patients hypertendus qui avaient des fonctions réservoir (31.23% \pm 9.93 vs 46.43% \pm 11.06, $p=0.000$) et de conduction de l'OG (14.26% \pm 2.91 vs 21.41% \pm 2.8, $p=0.000$) altérées par rapport aux patients normotendus. Au cours de la période contractile, le pic du strain de OG (16.73 \pm 3.84 vs 15.29 \pm 2.75, $p=0.07$) ainsi que le pic de sa vitesse étaient (-1.89% \pm 0.16, -1.82% \pm 0.21; $p=0.54$) plus élevés chez les patients hypertendus sans atteindre le seuil de significativité. Le temps d'atteinte du pic du strain de OG à la phase réservoir (405.02ms \pm 55.51 vs 387.13ms \pm 47.48, $p=0.05$) et la durée de la diastole (163ms \pm 26 vs 146ms \pm 24, $p=0.04$) étaient significativement plus élevés chez les patients hypertendus que chez les témoins. Une relation de corrélation significative entre les paramètres de l'étude volumétrique et ceux du speckle tracking a été notée.

Conclusion : Le strain longitudinal à la phase réservoir et conduite est altéré chez les patients ayant une hypertension artérielle malgré une normalité de la taille des cavités cardiaque et avant même l'altération des autres paramètres échocardiographiques conventionnels. Le speckle tracking est un outil prometteur de la détection des altérations précoces de l'OG

Mots Clés : Échocardiographie, Speckle Tracking, Oreillette Gauche, Hypertension, Fonction Diastolique, Fibrillation Auriculaire, Accident Vasculaire Cérébral Ischémique.

ABSTRACT

Introduction: Systemic hypertension is a major health problem worldwide, it is associated with impaired left atrial (LA) function. Myocardial deformation analysis using speckle-tracking echocardiography has emerged as a promising tool to evaluate atrial deformation and function.

Aim: To evaluate early changes in left atrial longitudinal strain based on speckle tracking echocardiography in patients with hypertension.

Methods: LA strain was studied using speckle-tracking echocardiography in 109 hypertensive patients without LA enlargement and 50 age-matched controls. Conventional and bidimensional strain echocardiographic assessments were performed and the following parameters were measured: peak atrial longitudinal strain and strain rate during the reservoir, conduit, and contractile periods in four and two-chambers views and time to peak atrial longitudinal strain/strain rate measured in the three phases of LA function.

Results: LA anteroposterior diameter was within the normal range, no difference between the hypertensive patients and controls was noted (34.35 mm \pm 4.91 vs 31.82 mm \pm 4.87, $p=0.16$). LA maximum volume (41.78ml \pm 10.29 vs 47ml \pm 13.21, $p=0.01$), minimum volume (23.95ml \pm 12.18 vs 16.94ml \pm 7.91, $p=0.001$) were higher in hypertensive patients, and impaired reservoir (31.23% \pm 9.93 vs 46.43% \pm 11.06, $p=0.000$) and conduit functions (14.26% \pm 2.91 vs 21.41% \pm 2.8, $p=0.000$) were noted in hypertensive patients compared to normotensive patients. During the contractile period, peak strain (16.73% \pm 3.84 vs 15.29% \pm 2.75, $p=0.07$) and strain rate (-1.89% \pm 0.16, -1.82% \pm 0.21; $p=0.54$) were higher in hypertensive patients without reaching the level of significance. Time to peak strain during reservoir period (405.02ms \pm 55.51 vs 387.13ms \pm 47.48, $p=0.05$) and duration of diastole (163ms \pm 26 vs 146ms \pm 24, $p=0.04$) were significantly higher in hypertensive patients compared to controls. A significant relationship between the parameters of the volumetric study and those of the bidimensional strain/strain rate study was noted.

Conclusion: Left atrial longitudinal strain during the reservoir and conduit periods is impaired in patients with hypertension despite normal cavity size and before the detection of other echocardiographic changes. Speckle-tracking echocardiography may be considered a promising tool for the early detection of LA strain abnormalities in these patients.

Keywords: Echocardiography, speckle tracking, left atrium, hypertension, diastolic function, atrial fibrillation, ischemic stroke

Correspondance

Karima Taamallah

Hôpital Militaire Principal d'Instruction de Tunis / Faculté de médecine de Tunis

Email: cardiokarima@yahoo.fr

INTRODUCTION

Hypertension (HTN) is a major health problem worldwide, and its prevalence is increasing. Unfortunately, its pathophysiology and clinical course are not yet clearly established. Several studies demonstrated a negative influence of HTN on target organ damage explaining the unfavorable prognosis of hypertensive patients and increased cardiovascular morbidity and mortality including atrial fibrillation, stroke, myocardial infarction, and sudden cardiac death in this population [1].

In hypertensive patients, comprehensive assessment of left atrial (LA) phasic function may be of clinical importance and might be helpful in the risk stratification in these patients. Indeed, LA enlargement and LA functional abnormalities may predict the occurrence of atrial fibrillation and cerebrovascular strokes in these patients [2,3]. Although structural and functional changes in the left ventricle during hypertension are well-known, relatively little is known about the effect of hypertension on LA functions and its prognostic impact, because they were insufficiently studied [3]. Some studies suggested that left atrial (LA) dysfunction and left ventricular (LV) diastolic dysfunction occur before structural changes of the LA and the LV even in patients with well-controlled HT.

Atrial structural and functional changes caused by hypertension can be evaluated with several conventional techniques such as echocardiography, computed tomography, magnetic resonance imaging. Two-dimensional (2DE) and three-dimensional (3DE) echocardiography are the most commonly employed noninvasive imaging techniques to evaluate LA size and function.

Speckle tracking echocardiography (STE) is an imaging technique that can be applied to the analysis of left atrial function. It allows direct and angle-independent analysis of myocardial deformation, thus providing sensitive and reproducible indices of myocardial fiber dysfunction that overcome most of the limitations of Doppler-derived strain measures, so, a more precise evaluation of the myocardium in patients with HTN, what is required to provide effective diagnosis and management of cardiac dysfunctions in patients with hypertension.

The assessment of LA function using bidimensional (2D) strain may be of particular interest in patients with no evidence of LA enlargement, because it may provide additional information for the early detection of LA abnormalities at the very early stage of the disease, which would improve clinicians' understanding and management of hypertensive patients and help to identify patients at high risk for adverse events especially heart failure

with preserved ejection fraction, and to predict atrial fibrillation (AF) and cardiac events. There have been few studies using STE to assess LA deformation in hypertensive patients and its prognostic impact [5].

So, we aimed in this current study to evaluate early changes in the left atrial function during the three mechanical LA phases by using two-dimensional speckle tracking echocardiography (2-DSTE) in hypertensive patients with normal LA size.

METHODS

We conducted a prospective cross-sectional study at our department of Cardiology between February 2015 and June 2020 including 159 subjects divided into two groups: The hypertensive group (HTN group) consisted of 109 patients with primary hypertension, and the control group comprised 50 healthy subjects. The two groups were age- and gender-matched and were comparable for most clinical variables. The study was conducted following the principles of the declaration of Helsinki and was approved by the local ethics committee. All participants gave informed consent to participate in the study.

Hypertension was defined as systolic blood pressure (BP) \geq 140 mmHg and/or diastolic BP \geq 90 mmHg or antihypertensive treatment with a documented history of hypertension. Blood pressure was measured before echocardiography, on the non-dominant arm by a sphygmomanometer in a sitting position according to current guidelines for the management of hypertension [1]. For each patient in both groups, anamnestic and clinical data were collected. Were included in this study, patients with an established diagnosis of hypertension and well-controlled by antihypertensive drugs, with normal BP (BP \leq 140 mmHg and/or diastolic BP \leq 90 mmHg) measured just before the practice of the echocardiography and with normal heart structure and functions: Normal LA size (LA anteroposterior diameter \leq 40 mm on the parasternal long-axis view), and volume (LA maximum volume index $<$ 34 ml/m²), normal left ventricular (LV) ejection fraction (EF) and without LV hypertrophy (LVH) (LVEF \geq 50% and normal LV mass $<$ 105g/ m² in male and $<$ 95g/ m² in female) (6).

Exclusion criteria

were excluded from this study, patients with conditions that affect LA size and function such as patients aged over 70 years, patients with secondary hypertension, with a history of ischaemic heart disease or angiographically documented coronary artery disease,

conduction abnormalities, sinus bradycardia and tachycardia, atrial and ventricular arrhythmias documented by electrocardiogram (ECG) or 24-h Holter ECG, permanent cardiac pacemaker implantation, bundle branch block, LV systolic dysfunction, and abnormal LV wall motion, aortic and mitral valvulopathy, previous cardiac surgery, diabetes mellitus, chronic obstructive pulmonary disease (COPD), obstructive pulmonary sleep apnea (OSA), and patients with poor echocardiographic window.

Echocardiography

All patients underwent an echocardiographic examination using the ultrasound machine General Electric Vivid9, with a 3.5-MHz transducer. The data were analyzed offline using EchoPAC (GE Vingmed Ultrasound AS). A standard echocardiographic study using 2D, M-mode, and Doppler techniques were performed in addition to speckle tracking for LA. All echocardiograms were performed by an experienced cardiologist, he was the same operator. The transthoracic echocardiography was performed with simultaneous recording of a stable and a good quality electrocardiogram trace. Cardiac dimensions and volumes were measured according to the American Society of Echocardiography's Guidelines [6].

Standard assessment of left atrium

LA diameter was derived from parasternal long-axis B-mode view, and LA volumes were calculated from apical four-chamber and two-chamber views using the biplane Simpson's method. All volumes were indexed to body surface area. LA maximum volume (LAVmax) was measured just before mitral valve opening, at the end of the T wave on systole, and LA minimum volume (LAVmin) was measured at mitral valve closure at the beginning of the QRS wave on the end of diastole. LA pre-atrial volume (LAVp) was measured just before the beginning of the P wave on the electrocardiogram.

The following parameters of the LA function were calculated

- Total LA stroke volume (LASV): Calculated as the difference between maximal and minimal LA volumes.
- LA expansion index (LAEI): Calculated as the ratio of total LASV to minimum LAVolume x 100.
- LA active emptying volume and LA passive emptying volume were calculated using (LAVp - LAVmin) and (LAVmax -LAVp) formulas, respectively.
- LA contractile function was assessed by calculating the LA active emptying fraction (LAAEF) that could be obtained as the ratio of LA active emptying volume to LAVp x 100.
- LA reservoir function was assessed by calculating LA ejection fraction (LAEF) that could be obtained as the ratio of LASV to LAVmax x 100.

- LA conduit function was assessed by calculating the LA passive emptying fraction (LApEF) that could be obtained as the ratio of LA passive emptying volume to LAVmax x 100. Left ventricular end-diastolic diameter (LVEDd), the interventricular septal (IVS), and LV posterior wall thickness in diastole (LVPW) in diastole were measured using parasternal long-axis view, left ventricular ejection fractions (LVEF) were calculated by Simpson's method. LV mass (LVM, in grams) was calculated using the Penn formula and was subsequently indexed to body surface area (BSA) to obtain LV mass index.

The classification of the LV geometry was based on the relative wall thickness. (RWT=2PW/EDD) and the LV mass : normal geometry (Normal LVM and a RWT < 0.42) and concentric remodelling (normal LVM and RWT> 0.42). Patients with LVH were excluded.

Diastolic parameters, including E/A and Em/Ea, were assessed using pulsed-wave Doppler at the tips of the mitral leaflets and from tissue Doppler imaging at the level of the lateral and septal annulus, respectively. The ratio of Em/Ea was calculated using the average Ea value.

Longitudinal Strain acquisition

Speckle tracking echocardiography recordings were processed using acoustic-tracking software (EchoPAC PC, Version113; GE Health VIVID 9), allowing off-line semi-automated analysis of speckle-based strain. Different strain parameters were acquired according to current recommendations [7].

Four and two apical-chambers views were acquired, using a conventional 2D gray-scale echocardiography, over three consecutive cardiac cycles at a rate of 70 images per second by asking the patient to perform an apnea for better image quality. A small sector angle (30°) was chosen to acquire the maximum frame rate, the software is applied in harmonic imaging. Care was taken to optimize visualization of the LA cavity and to maximize the LA area in apical views and avoid foreshortening of the left atrium. In the apical four-chamber and two-chamber views, the LA endocardial border was manually traced at end-systole in apical four-chamber and two-chamber views by a point-and-click approach. The system generated epicardial surface tracing automatically using the software automatically for each view. After manual adjustment of the region of interest (ROI) width and shape, the software divided the ROI into 6 segments for each view, and the resulting tracking quality for each segment was automatically scored as either acceptable or non-acceptable. Inadequately tracked segments were either corrected manually or rejected from the analysis.

The LA walls were divided into 12 segments, in subjects with

adequate image quality, a total of 12 segments were then analyzed. Lastly, the software generated the longitudinal strain curves for each segment and a mean curve of all segments that reflect the pathophysiology of atrial function. Setting zero strain at LV end-diastole, the LA strain pattern is characterized by a predominant positive wave that peaks at the end of ventricular systole, followed by two distinct descending phases in early diastole and late diastole. From apical four-chamber and two-chamber views, peak longitudinal LA strain (PS-S) and strain rate during ventricular systole (PSR-S) reflecting LA reservoir function was obtained just before the mitral valve opening. Peak strain (PS-E) and strain rate (PSR-E) are measured during early diastole and reflect LA conduit function during the conduit phase. Peak strain-A wave (PS-A) and peak strain rate-A wave (PSR-A) during late diastole reflecting pump function during LA contractile phase was obtained at the onset of the P-wave on electrocardiography. Peak atrial longitudinal strain during a different phase of LA function was calculated by averaging values obtained from all LA segments from apical four-chamber and two-chamber views. Figure 1 illustrates the LA strain mean curve of all LA segments and Figure 2 illustrates the LA strain rate mean curve of all LA segments. LA stiffness index (LASi) is calculated as the ratio of E_m/E_a to PS-S.

In the same manner, time to peak strain and strain rate were measured manually by using the time cursor of 2-DSTE. They were measured from the R wave of the electrocardiogram to the peak strain/strain rate during the reservoir, conduit, and contraction phases.

Duration of reservoir, conduit, and contractile phases were measured in hypertensive patients and controls, mean duration was calculated for each LA phase function.

Intraobserver and interobserver variability study

Intraobserver and interobserver variability were determined by repeating a Two-dimensional speckle-tracking study after three months for 30 randomly selected patients by the same observer and by a second independent observer.

Correlation study

The correlation between parameters measured by the conventional echocardiography and parameters measured by Speckle tracking imaging was studied. Absolute values of PSR-E and PSR-E were used for the determination of the correlation between these two parameters and conventional parameters.

Statistical study

SPSS for Windows software was used for statistical analysis (SPSS Inc. Chicago, USA). Peak strain and strain rate, SR, time to the peak strain, and strain rate of 12 LA wall during LA

relaxation, conduit, and contraction were compared between hypertensive and normotensive subjects. The homogeneity and normal distribution of continuous data were tested through the Leneve's test and Shapiro–Wilk test, respectively. A p-value of more than 0.05 was considered significant for the homogeneity and normal distribution values.

Categorical variables were expressed as absolute frequencies and percentages, Continuous non-parametric variables were expressed as the median value, and 25th–75th percentile and continuous parametric variables were expressed as the mean and standard deviation.

Categorical variables were compared by the likelihood-ratio χ^2 test. Comparison of parametric values between the two groups was performed by using a two-tailed Student's t-test. All values were obtained with 95% confidence intervals. A p-value of less than 0.05 was considered statistically significant.

Intraobserver and interobserver analyses were performed using Bland–Altman to evaluate the standard deviation and 95% limits of agreement.

Pearson's correlation coefficient (r) was used to determine a correlation between parameters of LA function from standard echocardiography and the 2D strain study.

RESULTS

General characteristics of the population

The mean age in the HTN group was 58.86 ± 11.61 years. The average duration of hypertension was 8.36 ± 6.37 years. As expected, systolic and diastolic blood pressure were significantly higher in the hypertensive group than those in the control group (table n°1).

Table 1. Clinical Characteristics of Hypertensive and Normotensive Subjects

	HTN (+) (n=109)	HTN (-) (n=50)	P-value
Age (years)	58.86 ± 11.61	54.84 ± 13.54	0.36
Men n (%)	58 (53%)	27 (54%)	0.49
Body surface area (m ²)	1.76 ± 0.2	1.74 ± 0.2	0.20
Body mass index (BMI) kg/m ²	27.2 ± 1.3	26.9 ± 1.1	0.82
Systolic blood pressure (mmHg)	149.5 ± 24.9	121.3 ± 10.5	<0.001
Diastolic blood pressure (mmHg)	90.1 ± 11.0	74.8 ± 6.7	<0.001
Heart rate (beats/min)	68.2 ± 8.8	62.4 ± 10.2	0.37
Duration of hypertension (years)	8.36 ± 6.37	-	-

Conventional echocardiographic characteristics

There was no difference between the two groups concerning LVEF, LVED, and LVES diameter, and ascending aortic diameter. Hypertensive patients had also greater septal and posterior wall thicknesses and LV mass, 69.7% (n=76) of patients have normal LV geometry. A significant difference was noted between the two groups for all diastolic function parameters (except Em /Ea ratio and DTm). Systolic pulmonary artery pressure (SPAP) was significantly increased in the hypertensive group (p=0.02). (Table 2).

Table 2. Conventional echocardiographic data of the two groups

	HTN (+)	HTN (-)	P-value
LV EDD (mm)	48.12± 5.94	46.76 ± 4.07	0.82
LV ESD (mm)	30.28 ±4.64	29.46 ±4.86	0.44
IVS (mm)	9.58±1.8	8.48±1.6	0.02
PW (mm)	9.10±1.53	8.16±1.31	0.05
LVM indexed to body surface area (g/m ²)	91.28±9.47	75.98±12.23	0.04
LVEF (%)	66.2 ±4.71	68.21 ±5.22	0.25
ascending aortic diameter (mm)	26.32±5.43	24.97±6.73	0.07
LA Diameter (mm)	34.35 ±4.91	31.82 ±4.87	0.16
LA area (cm ²)	16.22±3.66	13.96±2.74	0.02
LA maximum volume (ml)	41.78±10.29	47±13.21	0.01
LA maximum volume indexed to body surface area (ml/m ²)	28.60 ±5.4	27.5 ±6.17	0.01
LA minimum volume (ml)	23.95±12.18	16.94±7.91	0.001
LA minimum volume indexed to body surface area (ml/m ²)	14.38±7.22	10±4.71	0.001
LA stroke volume (ml/m ²)	20.36±4.21	24.34±2.18	0.001
LA stroke volume indexed to body surface area (ml/m ²)	14.3±6.1	15.9±5.8	0.02
LAVp (ml)	14.83±9.5 31	16.42±6.02	0.03
LAVp indexed to body surface area (ml/m ²)	26.41±10.42	28.4±3.8	0.04
LA ejection fraction (%)	62.53±9.12	66.8±1.10	0.05
LA expansion index (%)	1.7±0.83	2.0±0.9	0.03
LA stiffness index (%)	0.25±0.35	0.28±0.17	0.01
LApEF (%)	24.70±14.38	34.12±12	0.01
LAAEF (%)	48.65±17.28	44.10±13	0.06
Em (cm/s)	71.19±18.11	85.00±18.40	0.02
Am (cm/s)	85.56±18.23	66.83±16.77	0.001
Em/Am	0.94±0.33	1.35±0.46	0.000
DTm (ms)	218.53±57.68	205.00±47.52	0.06
Ea lateral (cm/s)	10.01±3.22	13.88±4.49	0.000
Ea average septal and lateral (cm/s)	9.86±2.47	11.74±3.08	0.049
Em /Ea	7.69±3.06	7.86±6.38	0.74
PSAP	29.85±5.11	23.07±08.74	0.02

LVEDD=left ventricular end diastolic diameter, LVESD= left ventricular end systolic diameter, IVS= interventricular septum, PW: posterior wall, LVM= left ventricular mass ; LVEF=Left ventricular ejection fraction ; LA= left atrium ; Em= transmitral Doppler early filling velocity;Am=indicates transmitral atrial filling velocity; DT= deceleration time, Ea= tissue Doppler early diastolic mitral annular velocity ; PSAP= pulmonary systolic arterial pressure, LAVp=left atrium passive emptying volume, LApEF= Left atrium passive emptying fraction, LAAEF= Left atrium active emptying fraction, HTN (+) : hypertensive patients, HTN(-) : Healthy controls, 4C= 4 chambers ; 2C= 2 chambers.

Left atrium volume and emptying fraction

Although all subjects in the hypertensive and control groups had normal LA size and volume, LA area, LAVmax, LAVmin, and LAVp were significantly higher in the hypertensive group than those in the control group (p<0.05). LA anteroposterior diameters were similar in the two groups. Hypertensive patients had lower LASV, LAEF, LAEI, LApEF, and LASi than controls (p<0.05). There was no difference between the two groups concerning LAAEF.

Left atrial 2D strain/strain rate

Left atrial 2D strain

In the hypertensive group, Peak strain (PS) measured in apical 4c and 2c views during reservoir phase (PS-S) and conduit phase (PS-E) were significantly lower with values of 31.23 ±9.93 versus 46.43 ±11.06 (p=0.000), and 14.26±2.91 % versus 21.41±2.8 % (p=0.000) respectively.

Although higher in hypertensive patients, there was no significant difference between the two groups regarding Peak strain obtained from both apical four-chamber and two-chamber views images during contractile phase (PS-A), with a value of 16.73 ±3.84% versus 15.29±2.75% (p=0.07). Comparison of LA longitudinal strain values between the two groups is shown in Table 3.

Table 3. Global longitudinal peak strain / strain rate values in the hypertensive group compared with the control group for the reservoir, conduit, and contractile phases.

	HTN (+)	HTN (-)	P-value
PS-S A4C (%)	30.22±9.96	45.16±10.11	0.000
PS-S A2C (%)	32.76±10.91	48.36±13.90	0.000
PS-S (%)	31.23 ±9.93	46.43 ±11.06	0.000
T-PS S (ms)	405.02±55.51	387.13±47.48	0.05
PS-E A4C (%)	14.69±2.79	21.13±3.4	0.000
PS-E A2C (%)	14.09±2.88	21.24±1.7	0.000
PS-E (%)	14.26±2.91	21.41±2.8	0.000
T-PS E (ms)	596±49	629±74	0.001
PS-AA4C (%)	16.46±3.12	15.31±2.75	0.08
PS-A A2C (%)	16.89±4.93	15.10±2.75	0.07
PS-A (%)	16.73 ±3.84	15.29±2.75	0.07
T PS-A(ms)	686±70	692±96	0.14
Reservoir phase duration (ms)	560±23	522±31	0.02
Conduit phase duration (ms)	68±13	71±16	0.06
Contraction phase duration (ms)	163±26	146±24	0.04
P-SR S A4c(%)	0.98±0.14	1.36±0.48	0.04
P-SR S A2c(%)	0.99±0.18	1.37±0.37	0.03
P-SR S(%)	0.99±0.12	1.37±0.19	0.03
P-SR E A4c(%)	-0.98±0.3	-1.58±0.6	0.001
P-SR E A2c(%)	-0.98±0.3	-1.51±0.7	0.002
P-SR E(%)	-0.98±0.3	-1.56±0.4	0.001
P-SR A A4c(%)	-1.87±0.19	-1.92±0.11	0.56
P-SR A A2c(%)	-1.9±0.24	-1.72±0.87	0.60
P-SR A(%)	-1.89±0.16	-1.82±0.21	0.54
T P-SR S(ms)	215±28	199±13	0.02
TP-SR E(ms)	498±11	436±24	0.01
TP-SR A(ms)	741±16	739±28	0.05

PS-S: Peak strain during reservoir phase, PS-E: Peak strain during conduit phase, PS-A: Peak strain during the contractile phase, T-PS S: Time to peak strain during reservoir phase, T-PS E: Time to peak strain during conduit phase, T-PS A: Time to peak strain during the contractile phase, A2C: Apical two chambers, A4C: Apical four chambers. LV: Left ventricular.

Left atrial 2D strain rate

Strain rate values were significantly impaired in the LA walls in hypertensive patients measured during reservoir (PSR-S) and conduit phases (PSR-E) from apical four-chamber et two-chamber views as well as their averages. The strain rate value measured during the contractile phase (PSR-A) was higher in hypertensive patients but this increase did not reach statistical significance (-1.89%±0.16 vs -1.82%±0.21,p=0,54). Global longitudinal peak strain rate values in the hypertensive group compared with the control group for reservoir, conduit, and contractile phases are shown in Table 3.

Duration of different phases of LA function

Compared to normal subjects, the mean duration of reservoir period (D-S) (560ms±23 , 522ms±31, p= 0.02) and contractile period (D-A) (163ms±26, 146ms±24 , p= 0.04) duration were significantly higher in hypertensive patients, but the mean duration of conduit period (D-E) (68 ms±13, 71ms±16;p= 71±16) was lower in hypertensive patients compared with healthy subjects, this decrease in D-E was not statistically significant (Table 3).

LA time to peak strain/strain rate

Mean LA time to peak strain/strain rate measured from both apical four-chamber and two-chamber views during the reservoir, conduit, and contractile phases were significantly higher in the hypertensive group except T-PSA which was higher in non-hypertensive patients without reaching the level of significativity (686±70, 692±96,p=0.14). (table 3).

Intra-and interobserver variability

Intra-and interobserver variability of PS-S measurements were 5.8% and 6.6%, respectively, of PS-E measurements were 7.2% and 8.7%, respectively, and of PS-A measurements were 7.4% and 7.9%, respectively.

Intra-and interobserver variability of PSR-S measurements were 5.3% and 7.6%, respectively, of PSR-E measurements were 6.2% and 8.1%, respectively, and of PSR-A measurements were 6.9% and 8.7%, respectively.

Correlation analysis between 2D strain and standard echocardiography parameters

No correlation was found between LA diameter and peak strain and strain rate in the reservoir, conduit, and contractile phases. LA area was correlated only with peak strain and strain rate in the contractile phase respectively (r=-0.143, p=0.01) (r=-0.143, p=0.01) (Table 4).

All echocardiographic parameters of the volumetric study, except LASV and LAAEF, correlated significantly with PS-S. A significant correlation was observed between PS-E and IVS, Em/Am, E/Ea, LAV max, LAV min, LApV, LASV, LAEI, LAEF, LApEF, and SPAP. LAVmin, LAV max, LAEF, and LAEI, were significantly correlated with PS-A, no statistically significant correlation was noted between PS-A and Em/Am, Em/Ea, LApV, LASV, LAEI, LApEF (see table 4).

Table 4. Correlations between standard echographic parameters and atrial 2d strain parameters

	PS-S	PS-E	PS-A	PSR-S	PSR-E	PSR-A
LA Diameter	r=-0.49 p=0.12	r=-0.49 p=0.16	r=-0.287 p=0.26	r=-0.49 p=0.18	r=-0.49 p=0.27	r=-0.287 p=0.34
LA area	r=-0.18 p=0.08	r=-0.08 p=0.09	r=-0.143 p=0.01	r=-0.13 p=0.12	r=-0.22 p=0.32	r=-0.143 p=0.01
LA maximum volume	r=-0.651 p=0.002	r=-0.64 p=0.002	r=-0.67 p=0.03	r=-0.54 p=0.04	r=-0.11 p=0.06	r=-0.12 p=0.42
LA minimum volume	r=-0.69 p=0.001	r=-0.67 p=0.002	r=-0.61 p=0.03	r=-0.013 p=0.32	r=-0.34 p=0.09	r=-0.53 p=0.01
Pre-atrial contraction volume	r=-0.64 p=0.002	r=-0.68 p=0.002	r=-0.15 p=0.30	r=-0.22 p=0.47	r=-0.48 p=0.04	r=-0.28 p=0.37
Stroke volume	r=0.15 p=0.19	r=0.43 p=0.01	r=0.13 p=0.12	r=0.41 p=0.04	r=0.18 p=0.06	r=0.28 p=0.34
LAEF	r=0.27 p=0.04	r=0.63 p=0.02	r=0.69 p=0.01	r=0.52 p=0.04	r=0.14 p=0.13	r=0.23 p=0.38
LAEI	r=0.62 p=0.00	r=0.52 p=0.03	r=0.38 p=0.00	r=0.21 p=0.06	r=0.19 p=0.07	r=0.12 p=0.52
LAAEF	r=0.21 p=0.06	r=0.16 p=0.07	r=0.12 p=0.6	r=0.31 p=0.07	r=0.24 p=0.08	r=0.63 p=0.001
LAEF	r=0.72 p=0.01	r=0.62 p=0.03	r=0.09 p=0.72	r=0.70 p=0.01	r=0.69 p=0.01	r=0.09 p=0.72
IVS	r=-0.62 p=0.02	r=-0.62 p=0.02	r=-0.39 p=0.19	r=-0.09 p=0.08	r=-0.15 p=0.14	r=-0.17 p=0.82
E/A	r=0.27 p=0.04	r=0.27 p=0.04	r=0.288 p=0.24	r=0.62 p=0.03	r=0.54 p=0.03	r=0.15 p=0.12
E/e'	r=-0.31 p=0.02	r=-0.31 p=0.03	r=0.28 p=0.35	r=-0.68 p=0.02	r=-0.61 p=0.02	r=-0.22 p=0.32
SPAP	r=-0.51 p=0.001	r=-0.34 p=0.001	r=-0.39 p=0.34	r=-0.53 p=0.01	r=-0.44 p=0.03	r=-0.17 p=0.26

LA: left atrium, LAEF: Left atrium ejection fraction, LAEI: Left atrium expansion index, LAAEF: Left atrium active emptying fraction, LApEF: Left atrium passive emptying fraction, IVS: inter-ventricular septum, SPAP: Systolic pulmonary artery pressure.

Correlation between 2D strain rate and standard echocardiography parameters

PSR-S was significantly correlated with LAVmax, LASV, LAEF, LApEF, Em/Am, Em/Ea, and SPAP (Table 4).

LAVp, LApEF, Em/Am, Em/Ea, and SPAP were correlated with PSR-E (Table 4).

Except for a correlation between PSR-A and LAAEF and LA minimum volume, no significant correlation existed between PSR-A and Em/Am, Em/Ea, LAV max, LApV, stroke volume, LAEI, LAEF, and LApEF (Table 4).

The correlation between standard echocardiographic parameters and 2D strain and strain rate study is shown in Table 4.

DISCUSSION

In our study, we used 2D strain and strain rate imaging to assess left atrial deformation in hypertensive patients with normal LA size. Despite normal LA size assessed by conventional 2D echocardiography, an impaired reservoir and conduit function in hypertensive patients compared to normotensive patients was noted. Besides, the correlation analysis indicated a significant relationship between parameters of the 2D strain/strain rate study and those of the volumetric study. During the contractile period, PS and PSR were higher in hypertensive patients without reaching the level of significance. Time to PS, T-PSR, and the duration of diastole were significantly higher in hypertensive patients compared to controls.

Physiopathology

Normal LA structure and function are essential for cardiac function with its three functions: Reservoir, conduct, and pump. Several studies suggest that LA structural remodeling and/or functional impairment might be involved in the pathogenesis and development of ventricular disorders, such as heart failure with reduced or preserved ejection fraction [8].

In hypertensive patients, complex morphological changes of LA and LV is the result of the heart adapting to the increased left ventricular workload. The LA enlargement is the result of forceful atrial contraction compensating for the reduction in early diastolic emptying due to diastolic dysfunction. Besides, the intermittent or permanent elevation of LV filling pressures leads to over LA filling leading to atrial fibrosis predisposing to atrial remodeling and dysfunction, and consequently to arrhythmia [9].

Speckle Tracking imaging

Advanced echocardiographic tools, such as two-dimensional (2DE) and, recently, three-dimensional (3DE) Speckle Tracking echocardiography, allow a better understanding of different cardiac chamber's function. It is used to measure both global and regional strain, defined as the degree of displacement of a region over the cardiac cycle, through tracking acoustic markers generated by the effect of ultrasound on the myocardium. Clinical implications of STE are various, it has shown to be sensitive for the detection of subclinical disease, including hypertensive heart disease [2]. It was suggested that speckle-tracking echocardiography is more accurate than LA size or volume to detect an early LA dysfunction [10-12].

The interest of speckle tracking in hypertensive patients

Our findings extend previous reports describing LA deformation during the three phases of LA function.

Several studies have shown that abnormal LA deformation precedes LA enlargement and LV hypertrophy. It was demonstrated that in hypertensive patients with normal LA size, peak strain, and strain rate were significantly lower during reservoir and conduit phases [13-16].

An increased LASi was proposed by some authors as a marker of early target organ damage in hypertension [16], this parameter is decreased in our patients.

The influence of left ventricular geometry on the left atrial phasic function was reported by several authors [16-19]. The decrease in reservoir and conduct functions are more pronounced in hypertensive patients with concentric LV than in hypertensive patients with normal geometry or concentric remodeling. Several studies have demonstrated that the decrease of different indices of LA function was significantly pronounced in the hypertensive patients in the presence of diabetes mellitus, a higher grade of LV diastolic dysfunction, higher age, and obesity [5,20-24].

In opposition to our results, an impairment of atrial function during the contraction phase in addition to the decreased strain during reservoir and conduit has been demonstrated by several authors [9, 25].

Dernellis et al. [26] have demonstrated an increase in reservoir and pump booster function which was in contrast with our results, in this study conduit function was decreased, the alteration of LA conduit function would be related to the increase in LV filling pressure during diastole.

In our study, strain and strain rate parameters were increased during the contraction phase in hypertensive patients

without reaching the level of significance, a significant increase in LA booster pump function was revealed by several studies [9,11,27,28]. It was suggested that LA function during atrial contraction is a potential non-invasive indicator of left ventricular end-diastolic compliance [7]. According to some authors, the increase in LA contraction function was pronounced in patients with concentric LVH compared to subjects with normal LV geometry [17].

Several studies suggest that the decreased LA strain is a good sign of LV diastolic dysfunction, they have shown that there is a good correlation between PS-S and LV diastolic function as well as increased LV filling pressures. According to some authors, LA strain measurements allow accurate diagnosis and categorization of diastolic dysfunction. Since LA conduit and reservoir functions decrease earlier than the diagnosis of LV diastolic dysfunction, some authors have proposed these parameters in the decisional algorithm to study filling pressures [28, 29].

Left atrial dysfunction was also demonstrated in patients with masked hypertension. Soe M. Aung et al. [30] have demonstrated the utility of 2D strain to detect masked hypertension in patients with hypertensive response to exercise. A decreased reservoir and pump function in these patients was noted. The same results were demonstrated by Tadic et al. [31], who suggest that 24-hour systolic blood pressure increment was closely related to LA remodeling.

A disturbance in LA function was also noted in patients with white coat hypertension. Tadic et al. [32] have demonstrated that patients with white coat hypertension have a reduced conduit function and an increased pump function, however, reservoir function was normal. These authors have compared LA function in patients with sustained hypertension, white coat hypertension, and controls, the LA function changes mentioned above, were more pronounced in sustained hypertensive patients. In this same study, LA stiffness, as well as aortic stiffness, gradually increases from controls to sustained hypertensives.

A Lower peak strain and strain rate during reservoir, conduit, and contraction phase have been reported in hypertensive patients with obstructive sleep apnea (OSA) compared to hypertensive patients without OSA, highlighting, according to some authors, the potential role of OSA in increasing the risk of atrial fibrillation in these patients [33].

In patients with non-dipper hypertension, LA function is

more impaired compared to dipper hypertensive patients [34-37]. It was demonstrated that the raised LV pressure secondary to the nocturnal systemic pressure overload was strongly associated with LA deformation [38].

A decrease in LA strain was also found in cases of gestational hypertension. Andrea Sonaglioni et al. [5] reported an impaired LA function in women with new-onset gestational hypertension, these authors have demonstrated an incremental prognostic value of global left atrial peak strain in these patients.

Gee Hee Kim et al. [38] revealed that compared with normotensive subjects, never treated early hypertensive patients and without clinically apparent target organ damage, had significantly increased plasma aldosterone concentration, and decreased atrial reservoir pump strain, and atrial systolic strain rate. LA pump strain was independently associated with nighttime systolic BP. Plasma aldosterone concentration was correlated with LV deformation. No correlation was found between LA deformation and plasma aldosterone level. Recently, in hypertensive and diabetic patients, Kalaycıoğlu et al. [39] have demonstrated that higher serum osteoprotegerin level was associated with impaired LA function assessed by speckle tracking. These authors suggested that serum osteoprotegerin may be used as a risk predictor for LA mechanical dysfunction.

Speckle tracking echocardiography is useful as a monitoring tool predicting the response to hypertension treatment and the reversibility of structural anomalies. Dernellis et al. [26] have demonstrated an improvement in atrial function in hypertensive patients who had adequate hypotensive treatment by ACE inhibitors and/ or thiazide diuretics. According to these authors, this improvement in atrial function might be explained by the regression of LV hypertrophy and the contribution of the autonomic nervous system. Ya suo et al. [40] have demonstrated that treatment by ACE inhibitors is associated with reduced risk of left atrial appendage thrombosis formation in hypertensive patients with atrial fibrillation. Several other studies have demonstrated a beneficial effect of ACE on left atrial function [19, 41].

In patients with suboptimal blood pressure control, Chen XJ et al. [42] have demonstrated a decreased left atrial myocardial strain. These authors suggest that suboptimal BP control status in hypertensive patients is related to a further reduction

of LA myocardial reservoir, conduit, and pump function, they thought that suboptimal BP might be regarded as a composite risk factor and therefore a simplified treatment target.

Warita et al. [43] have demonstrated that Pitavastatin had a beneficial effect on LV diastolic function and LA structure and function in elderly patients with HTN. Pitavastatin treatment may be associated with a lower incidence of new-onset atrial fibrillation [AF].

From a prognostic point of view, it is widely known that hypertension is a risk factor for developing AF [44]. In addition to LA enlargement, LA functional abnormalities may also predict the occurrence of atrial fibrillation [45]. According to some authors, LA strain parameters could be useful predictors of AF occurrence in hypertensive patients [46,47]. So, in patients with impaired atrial function, closer surveillance to detect arrhythmia is necessary. Table 5 summarizes data of different published studies on left atrium deformation assessment in hypertensive patients.

Table 5. Data of different published studies on left atrium deformation assessment in hypertensive patients

Authors	Year	Software	The population of the study n patients	Reservoir Function S (%) SR (L/s)	Conduit Function S (%) SR (L/s)	Pump Function S (%) SR (L/s)
T.XU et al.(9)	2015	E9, General Electric	124 HTN	45.3±7.7 2.3±0.5	20.2±3.8 2.2±0.6	25.0±5.9 2.6±0.5
Sahebjam et al.(13)	2014	7 Vivid 7, GE Healthcare	75 HTN	14.98±5.86 1.31±0.5	- -	- -
Liu et al. (14)	2014	Vivid 7 (GE Vingmed Ultrasound, Horten, Norway) equipped	99HTN and 65 HTN +diabetes	29.2±7.9 1.2±0.3	14.9±5.5 1.1±0.4	14.3±4.1 1.4±0.5
Karakurt et al.(15)	2019	Vivid 7, GE Healthcare	55 HTN	11.25±5.81 1.22±0.45	2.69±2.79 -0.95±0.54	11.63±6.2 -1.62±0.8
Kukubu et al. (19)	2007	-	55 HTN without LA dilation	2.15±0.57		
Hennawy et al.(24)	2018	Vivid S5 or Vivid 9; GE Medical Systems. A	50 HTN	24±6.29 -	- -	- -
Soullier et al.(25)	2016	Vivid 9 General electric medical health Echo PAC software	30 HTN with LVH	23±6 1.11±0.4	11.3±5 0.91±0.4	12±4 1.47±0.43
Miyoshi et al. (28)	2013	Vivid 7, General Electric Healthcare, Milwaukee, WI	126 HTN	31.4±8.9 1.4±0.4	16.6±6.2 -1.3±0.5	14.8±4.8 -1.7±0.6
Medeiros et al.(34)	2020	CX-50 Philips Healthcare	65 HTN	51.2 2.1±0.4	30.9±5.7 -2.89±0.7	20.1±5.6 -3.12±0.7
Gee Hee Kim et al. (39)	2019	Vivid E9, GE Medical Systems, Horten, Norway	71 never treated hypertensive patients	33.2 ±7 11.42 ± 0.31	18.2 ± 6.3 -1.51 ± 0.54	14.9 ± 3.8
Dimitroula et al.(42)	2010	-	40 HTN	63.3±4.1 3.1 ±0.2	- -	- -
Chen et al.(43)	2017	IE 33 Philips med system Qlab 7.0	279 HTN	35.9±8 -	18.5±7.1 -	17.8±4.2 -
Watanabe et al.(48)	2013	ACUSON sequoia 512 (Siemens, Mountain View, CA, USA) Normotensive	163 HTN	25.5±0.92 1.18±0.34	13.68±0.59 -1.02±0.31	- -1.38±0.42
Our study		Vivid E9, GE Medical Systems, Horten, Norway	109 HTN with normal LA size and function	31.23 ±9.93 0.99±0.12	14.26±2.91 -0.98±0.3	16.73 ±3.84 -1.89±0.16

HTN: hypertension, LVH: left ventricle hypertrophy, LA: Left atrium

Limitations

The measurement of LA 2D strain is feasible and reproducible in most cases [48], but relies very widely upon operator skills and adequate apical views. This modality of echocardiography is less accurate in patients with non-sinus rhythm, requiring the average value of almost five consecutive beats. The disadvantages of LA 2D strain are also the possibility of error in the tracing of endocardial contours and the strict dependence of the image rate. The other pitfall of this technique is that the analysis is performed on the left ventricular strain software because there is no software for atrial strain [49, 50].

Limitations of the present study are the small sample, explained by the exclusion of patients with diabetes mellitus and/or another disease that may lead to LA dysfunction, and patients with AF, who constitute a significant percentage of hypertensive patients, and the number of patients who had to be excluded because of inadequate image quality for measuring LA 2D strain that is angle independent technique but necessitates a good quality imaging.

It would be interesting to conduct further studies with larger populations and follow-up data to precise the discriminatory role of LA 2D strain in the AF and heart failure with preserved ejection fraction risk stratification. Indeed, identifying hypertensive patients at risk for developing these complications by early diagnosis of LA dysfunction, and prompt institution of effective treatment should be the goal when considering this patient population.

CONCLUSION

Our study demonstrated that speckle tracking imaging could be used to detect subtle impairment of LA function in patients with hypertension. Hypertension is associated with impaired reservoir and conduit LA function and higher booster function, which may be compensatory. The clinical usefulness of LA function by STE in these patients merits further investigations to better precision of the role of the LA study in the prediction of atrial fibrillation, and the risk of heart failure with preserved ejection fraction.

REFERENCES

- Williams B, Mancia G, Spiering W, *and al.* 2018 practice guidelines for the management of arterial hypertension of the European society of cardiology and the European society of hypertension ESC/ESH task force for the management of arterial hypertension. *Journal of hypertension* 2018; 36, 2284-2309.
- Rijcken J, Bovenoeerd PHM, Schoofs AJG, Van Campen DH, Arts T. Optimization of cardiac fiber orientation for homogeneous fiber strain during ejection. *Ann Biomed Eng* 1999; 27, 289-297.
- Roşca M, Lancellotti P, Popescu BA, Piérard LA. Left atrial function: Pathophysiology, echocardiographic assessment, and clinical applications. *Heart* 2011; 97,1982-1989.
- Mondillo S, Cameli M, Caputo ML, *and al.* Early detection of left atrial strain abnormalities by speckle-tracking in hypertensive and diabetic patients with normal left atrial size. *J Am Soc Echocardiogr* 2011 ; 24, 898-908.
- Sonaglioni A, Lonati C, Lombardo M, *and al.* Incremental prognostic value of global left atrial peak strain in women with new-onset gestational hypertension. *J Hypertens* 2019; 37,1668-1675.
- Lang RM, Badano LP, Victor MA, *et al.* Recommendations for cardiac chamber quantification by echocardiography in adults: An update from the American Society of Echocardiography and the European Association of Cardiovascular Imaging. *J Am Soc Echocardiogr* 2015; 28, 1-14.
- Badano LP, Koliaş TJ, Muraru D, *and al.* Standardization of left atrial, right ventricular, and right atrial deformation imaging using two-dimensional speckle tracking echocardiography: A consensus document of the EACVI/ASE/Industry Task Force to standardize deformation imaging. *Eur Heart J Cardiovasc Imaging* 2018;19, 591-600.
- Xu TY, Sun JP, Lee APW, *and al.* Left atrial function as assessed by speckle-tracking echocardiography in hypertension. *Med* 2015; 94, 1-7.
- Bello V Di, Talini E, Dell'Omo G, *and al.* Early left ventricular mechanics abnormalities in prehypertension: A two-dimensional strain echocardiography study. *Am J Hypertens* 2010; 23, 405-412.
- Onishi N, Kawasaki M, Tanaka R *and al.* Comparison between left atrial features in well-controlled hypertensive patients and normal subjects assessed by three-dimensional speckle tracking echocardiography. *J Cardiol* 2014;63:291-5.
- Furukawa A, Ishii K, Hyodo E *and al.* Three-dimensional speckle tracking imaging for assessing left atrial function in hypertensive patients with paroxysmal atrial fibrillation. *Int Heart J* 2016; 57:705-11.
- Sahebjam M, Mazareei A, Lotfi-Tokaldany M, Ghaffari N, Zoroufian A, Sheikhatollahi M. Comparison of Left Atrial Function between Hypertensive Patients with Normal Atrial Size and Normotensive Subjects Using Strain Rate Imaging Technique. *Arch Cardiovasc Imaging* 2014; 2, 1-6.
- Liu Y, Wang K, Su D *and al.* Noninvasive assessment of left atrial phasic function in patients with hypertension and diabetes using two-dimensional speckle tracking and volumetric parameters. *Echocardiography* 2014; 31, 727-735.
- Karakurt A, Yildiz C, Yildiz A *and al.* Early detection strain/strain rate and time to strain/strain rate abnormalities for left atrial mechanical function in hypertensive patients. *Acta Cardiol* 2019;74,141-51.
- Yang Y, Zhang BW, Qi LT, Ma W, Meng L HY. Speckle tracking analysis of left atrial phasic function in patients with

- hypertension. *Journal Of Pekin University* 2014;46,596–600.
16. Zhao Y, Sun Q, Han J, *and al.* Left atrial stiffness index as a marker of early target organ damage in hypertension. *Hypertens Res* 2020;44:299-309.
 17. Eshoo S, Semsarian C, Ross DL, Marwick TH, Thomas L. Comparison of left atrial phasic function in hypertrophic cardiomyopathy versus systemic hypertension using strain rate imaging. *Am. J. Cardiol* 201;107,290–6.
 18. Kokubu N, Yuda S, Tsuchihashi K *and al.* Noninvasive assessment of left atrial function by strain rate imaging in patients with hypertension: a possible beneficial effect of renin-angiotensin system inhibition on left atrial function. *Hypertens Res* 2007;30:13–21.
 19. Tadic M, Cuspidi C, Ilic I *and al.* The relationship between blood pressure variability, obesity and left atrial phasic function in hypertensive population. *Int J Cardiovasc Imaging* 2016;32:603–12.
 20. Miyoshi H, Oishi Y, Mizuguchi Y *and al.* Contribution of obesity to left atrial and left ventricular dysfunction in asymptomatic patients with hypertension: A two-dimensional speckle-tracking echocardiographic study. *J Am Soc. Hypertens* 2014.8 :54–63.
 21. Ikejder Y, Sebbani M, Hendy I, Khramz M, Khatouri A, Bendriss L. Impact of Arterial Hypertension on Left Atrial Size and Function. *Biomed Res Int* 2020; 4: 1-7.
 22. Yang Y, Zhang B, Qi L, Ma W, Meng L HY. Synergistic effect of hypertension and aging on left atrial volume and function. *Zhonghua Nei Ke Za Zhi* 2014; 53:617–21.
 23. Hennawy B, El Kilany W, Galal H, Mamdouh A. Role of speckle tracking echocardiography in detecting early left atrial dysfunction in hypertensive patients. *Egypt Heart J* 2018;70, 217–23.
 24. Soullier C, Niamkey JT, Ricci JE, Messner-Pellenc P, Brunet X, Schuster I. Hypertensive patients with left ventricular hypertrophy have global left atrial dysfunction and impaired atrio-ventricular coupling. *J Hypertens* 2016; 34 1615–1620.
 25. Erol MK, Yilmaz M, Acikel M KS. Left atrial mechanical function in patients with essential hypertension. *Acta Cardiol* 2002; 57: 323–7.
 26. Dernelis JM, Vyssoulis GP, Zacharoulis AA TP. Effects of antihypertensive therapy on left atrial function. *J Hum Hypertens* 1996;10:789–94.
 27. Miyoshi H, Oishi Y, Mizuguchi Y *and al.* Early predictors of alterations in left atrial structure and function related to left ventricular dysfunction in asymptomatic patients with hypertension. *J Am Soc Hypertens* 2013;7:206–15.
 28. Oki T, Miyoshi H, Oishi Y, Mizuguchi Y, Ara N IA. The Impact of Hypertension as a Road to Heart Failure with Preserved Ejection Fraction: Diagnostic Value of Two-Dimensional Speckle Tracking Echocardiography for the Early Impairment of Left Atrial-Left Ventricular-Arterial Coupling. *Curr Hypertens Rev* 2014;10:177-88.
 29. Jarasunas J, Aidietis A, Aidietiene S. Left atrial strain - An early marker of left ventricular diastolic dysfunction in patients with hypertension and paroxysmal atrial fibrillation 11 Medical and Health Sciences 1102 Cardiorespiratory Medicine and Haematology. *Cardiovasc Ultrasound* 2018;16:1–9.
 30. Aung SM, Güler A, Güler Y, Karabay CY, Akdemir I. Two-dimensional speckle-tracking echocardiography-based left atrial strain parameters predict masked hypertension in patients with hypertensive response to exercise. *Blood Press Monit* 2017 ; 22 :27–33.
 31. Tadic M, Cuspidi C, Radojkovic J, Rihor B, Kocijanic V, Celic V. Masked Hypertension and Left Atrial Dysfunction: A Hidden Association. *J Clin Hypertens* 2017;19:305–311.
 32. Tadic M, Cuspidi C, Pencic B, *and al.* The influence of white-coat hypertension on left atrial phasic function. *Blood Press* 2017;26:102–108.
 33. Medeiros MA, Lins-Filho OL, Patriota TLGC, *and al.* Abnormal atrial function in hypertensive patients with obstructive sleep apnea assessed by speckle tracking echocardiography. *Hypertens Res* 2020; 43: 841–844.
 34. Açar G, Bulut M, Arslan K *and al.* Comparison of left atrial mechanical function in nondipper versus dipper hypertensive patients: A speckle tracking study. *Echocardiography* 2013;30:164–70.
 35. Demir M, Aktaş İ, Yıldırım A. Left atrial mechanical function and stiffness in patients with nondipper hypertension: A speckle tracking study. *Clin Exp Hypertens* 2017;39:319–24.
 36. Tosun V, Korucuk N, Kiliç AY *and al.* Evaluation of atrial electromechanical functions in dipper and nondipper hypertension patients using left atrial strain P-wave dispersion and P terminal force. *Echocardiography* 2018 ;35:1318–25.
 37. Tadic M, Cuspidi C, Majstorovic A *and al.* Does a nondipping pattern influence left ventricular and left atrial mechanics in hypertensive patients? *J. Hypertens* 2013 ;31:2438–46.
 38. Kim GH, Cho JS, Chung WB, Park CS, Ihm SH. Association of the plasma aldosterone concentration, left atrial deformation and ambulatory blood pressure in never-treated early hypertensive patients. *Clin Exp Hypertens* 2019;41: 651–656.
 39. Kalaycioğlu E, Çetin M, Çinier G, *and al.* Serum osteoprotegerin level is independently related to subclinical left atrial mechanical function in patients with hypertension and diabetes. *Herz* 2020;5:1-200.
 40. Suo Y, Zhang Y, Wang Y, *and al.* Renin–Angiotensin system inhibition is associated with reduced risk of left atrial appendage thrombosis formation in patients with atrial fibrillation. *Cardiol* 2018 ;25 :611–620.
 41. Dimitroula H, Damvopoulou E, Giannakoulas G *and al.* Effects of renin-angiotensin system inhibition on left atrial function of hypertensive patients: an echocardiographic tissue deformation imaging study. *Am J Hypertens* 2010; 23:556-61.
 42. Chen XJ, Chen C, Liang YJ, *and al.* Decreased left atrial myocardial strain in patients with suboptimal blood pressure control. *Clin Exp Hypertens* 2017 ;39: 481–488.
 43. Warita S, Kawasaki M, Tanaka R, *and al.* Effects of pitavastatin on cardiac structure and function and on prevention of atrial fibrillation in elderly hypertensive patients: A prospective study of 2-years' follow-up. *Circ J* 2012; 76: 2755–2762.
 44. Cuspidi C, Meani S, Fusi V, Valerio C, Catini E, Sala C, Sampieri L, Magrini F ZA. Prevalence and correlates of left atrial enlargement in essential hypertension: role of ventricular geometry and the metabolic syndrome: The Evaluation of Target Organ Damage in Hypertension study.

J Hypertens 2005 ;23 :875–82.

45. Tsang TS, Barnes ME, Bailey KR, Leibson CL, Montgomery SC, Takemoto Y, Diamond PM, Marra MA, Gersh BJ, Wiebers DO, Petty GW SJ. Left atrial volume: important risk marker of incident atrial fibrillation in 1655 older men and women. *Mayo Clin Proc* 2001 ;76: 467–75.
46. Petre I, Onciul S, Iancovici S and al. Left Atrial Strain for Predicting Atrial Fibrillation Onset in Hypertensive Patients. *High Blood Press Cardiovasc Prev* 2019; 26 :331–7.
47. Watanabe T, Kawasaki M, Tanaka R and al. Association among blood pressure control in elderly patients with hypertension, left atrial structure and function and new-onset atrial fibrillation: A prospective 2-year study in 234 patients. *Hypertens Res* 2013; 36 :799–806.
48. Rausch K, Shiino K, Putrino A, Lam AK yin, Scalia GM, Chan J. Reproducibility of global left atrial strain and strain rate between novice and expert using multi-vendor analysis software. *Int J Cardiovasc Imaging* 2019;35:419–426.
49. Cameli M, Ciccone MM, Maiello M, and al. Speckle tracking analysis: A new tool for left atrial function analysis in systemic hypertension: An overview. *J Cardiovasc Med* 2016;17:339–343.
50. Voigt JU, Malaescu GG, Haugaa K, Badano L. How to do la strain. *Eur Heart J Cardiovasc Imaging* 2020; 21: 715–717.